

AD-A078 487

BROWN UNIV PROVIDENCE R I LEFSCHETZ CENTER FOR DYNAM--ETC F/6 12/1
OPTIMAL DISCOUNTED LINEAR CONTROL OF THE WIENER PROCESS.(U)
SEP 79 I KARATZAS

AFOSR-77-3063

UNCLASSIFIED


AFOSR-TR-79-1249

NL

1 OF 1
AD-A
A 078487



END
DATE
FILMED
1-80
DDC



NATIONAL BUREAU OF STANDARDS
MICROCOPY RESOLUTION TEST CHART

UNCLASSIFIED ~~CONFIDENTIAL~~ ~~SECRET~~

LEVEL II

18 REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM	
1. REPORT NUMBER AFOSR-TR-79-1249		3. RECIPIENT'S CATALOG NUMBER	
4. TITLE (and Subtitle) OPTIMAL DISCOUNTED LINEAR CONTROL OF THE WIENER PROCESS		5. TYPE OF REPORT & PERIOD COVERED Interim rept.	
7. AUTHOR(s) Ioannis Karatzas		6. PERFORMING ORG. REPORT NUMBER	
9. PERFORMING ORGANIZATION NAME AND ADDRESS Brown University Division of Applied Mathematics Providence, RI 02912		8. CONTRACT OR GRANT NUMBER(s) AFOSR-77-3063	
11. CONTROLLING OFFICE NAME AND ADDRESS Air Force Office of Scientific Research/NM Bolling AFB, Washington, DC 20332		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 61102F 2304/A1 17A1	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE September 1979	
		13. NUMBER OF PAGES 16	
		15. SECURITY CLASS. (of this report) UNCLASSIFIED	
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE	
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)			
18. SUPPLEMENTARY NOTES			
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)			
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) The following stochastic control problem is considered: to minimize the discounted expected total cost $J(x;u) = E \int_0^\infty e^{-\alpha t} [\phi(x_t) + u_t(x)] dt$ subject to $dx_t = u_t(x)dt + dw_t$, $x_0 = x$; $ u_t \leq 1$, (w_t) a			

DDC
RECEIVED
DEC 20 1979
B

DDC FILE COPY

AD A 078487

DD FORM 1 JAN 73 1473

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

401 834

20. Abstract continued.

Wiener process, $\alpha > 0$. All bounded by unity, measurable and nonanticipative functionals $u_t(x)$ of the state process (x_t) are admissible as controls. It is proved that the optimal law is of the form

$$\begin{aligned} u_t^*(x) &= -1, & x_t &> b \\ &= 0, & |x_t| &\leq b \\ &= 1, & x_t &\leq -b \end{aligned}$$

for some switching point $b > 0$, characterized in terms of the function $\phi(\cdot)$ through a transcendental equation.

ACCESSION for	
NTIS	White Section <input checked="" type="checkbox"/>
DDC	Buff Section <input type="checkbox"/>
UNANNOUNCED	<input type="checkbox"/>
JUSTIFICATION	
BY	
DISTRIBUTION/AVAILABILITY CODES	
Dist.	AVAIL. and/or SPECIAL
A	

OPTIMAL DISCOUNTED LINEAR CONTROL OF THE
WIENER PROCESS^{*}

by

Ioannis Karatzas
Lefschetz Center for Dynamical Systems
Division of Applied Mathematics
Brown University
Providence, Rhode Island 02912

September 1979

^{*}This research was supported by the Air Force Office of Scientific Research under AF-AFOSR 77-3063.

OPTIMAL DISCOUNTED LINEAR CONTROL OF THE
WIENER PROCESS⁺

Ioannis Karatzas

ABSTRACT

The following stochastic control problem is considered: to minimize the discounted expected total cost

$$J(x;u) = E \int_0^{\infty} e^{-\alpha t} [\phi(x_t) + |u_t(x)|] dt$$

subject to $dx_t = u_t(x)dt + dw_t$, $x_0 = x$; $|u_t| \leq 1$, (w_t) a Wiener process, $\alpha > 0$. All bounded by unity, measurable and nonanticipative functionals $u_t(x)$ of the state process (x_t) are admissible as controls. It is proved that the optimal law is of the form

$$\begin{aligned} u_t^*(x) &= -1, & x_t &> b \\ &= 0, & |x_t| &\leq b \\ &= 1, & x_t &< -b \end{aligned}$$

for some switching point $b > 0$, characterized in terms of the function $\phi(\cdot)$ through a transcendental equation.

AIR FORCE OFFICE OF SCIENTIFIC RESEARCH (AFSC)
NOTICE OF TRANSMITTAL TO DDC

This technical report has been reviewed and is approved for public release IAW AFR 100-12 (7b). Distribution is unlimited.

A. D. BLOSE
Technical Information Officer

1. INTRODUCTION AND SUMMARY

→ We consider the problem of discounted optimal control of the Wiener process $\{w_t; t \geq 0\}$ transformed by the action of a non-anticipative control into the "state process" $\{x_t; t \geq 0\}$. The latter satisfies the "state equation" $dx_t = u_t(x)dt + dw_t, t \geq 0; x_0$ on an appropriate probability space.

→ There is a cost $\phi(x_t)$ per unit time for being in the wrong state x_t ^{which} where $\phi(\cdot)$ is an even, uniformly convex function on the reals whose second derivative is decreasing with distance from the origin. There is also a cost $|u_t|$ per unit time for using the control, u_t . Both costs are discounted in time by the factor $e^{-at}, a > 0$ and the control action is limited, $|u_t| \leq 1, a.s.$

→ The controller has to choose a law $u_t(x)$ as a nonanticipative, measurable functional of the state process with values in $[-1,1]$, so as to minimize the expected discounted total cost. → next page

The "physically obvious" law is to push with full force (to the right direction) if x_t is outside a certain neighbourhood of the origin, while to exert no control at all if x_t is in this neighbourhood:

$$(1.1) \quad \begin{aligned} u_t^*(x) &= -1, & x_t &> b \\ &= 0, & |x_t| &\leq b \\ &= 1, & x_t &< -b \end{aligned}$$

Optimality of this law is proved and the cutoff point b separating the active region from the dead zone is characterized in terms of the function $\phi(\cdot)$ through the transcendental

equation (3.11). Existence and uniqueness of a solution to the above equation is proved by making use of the aforementioned properties of $\phi(\cdot)$. It is an interesting problem to relax these assumptions in order to allow cost functions with general polynomial or even exponential growth.

General existence results for the problem of discounted stochastic control were given by Kushner [1967]. Beneš, Shepp and Witsenhausen [1979] proved optimality of the bang-bang law in the case of a quadratic running cost on the state and no cost on the control. They also treated the finite-fuel problem with a discounted cost criterion.

^{cont} In the present paper we proceed by formulating the control problem, in ~~Section 2.~~ The Bellman equation of dynamic programming is explicitly solved, in ~~Section 3.~~ and the candidate for the optimal law ~~u^* in (1.1)~~ is discerned from the properties of the solution. Optimality of the candidate is proved, in ~~Section 4.~~

KEY WORDS AND PHRASES: Discounted stochastic control, Bellman equation, dead-zone controllers

2. THE STOCHASTIC CONTROL PROBLEM

Consider as basic probability space Ω the space $C(\mathbb{R}^+)$ of continuous, real-valued functions on \mathbb{R}^+ and let \mathcal{F}_t , $t \geq 0$ denote the σ -field generated by $\{x_s; s \leq t\}$, $x \in \Omega$. Consider also the σ -field \mathcal{A} generated by the subsets M of $\mathbb{R}^+ \times C(\mathbb{R}^+)$ with the property that each t -section M_t of M belongs to \mathcal{F}_t and each x -section M_x of M is Lebesgue measurable. A function g defined on $\mathbb{R}^+ \times C(\mathbb{R}^+)$ is \mathcal{A} -measurable if and only if $g(t, \cdot)$ is \mathcal{F}_t -measurable for any $t \geq 0$ and $g(\cdot, x)$ is Lebesgue measurable, for any $x \in C(\mathbb{R}^+)$.

Definition 2.1: Let the control measure space be the compact interval $[-1, 1]$ with its Borel sets. An admissible nonanticipative control u is a measurable function $u: (\mathbb{R}^+ \times C(\mathbb{R}^+)) \rightarrow [-1, 1]$.

The class of all such controls is denoted by \mathcal{U} . For any control law $u \in \mathcal{U}$ and any $x \in \mathbb{R}$ we can construct by means of the Girsanov theorem a probability space (Ω, \mathcal{F}, P) and a pair of stochastic processes (x_t, w_t) on it, such that $\{w_t; t \geq 0\}$ is a Wiener process with respect to P and the stochastic differential equation

$$(2.1) \quad dx_t = u_t(x)dt + dw_t, \quad t \geq 0$$

$$(2.2) \quad x_0 = x$$

is satisfied. Such a "weak solution" of (2.1) is known to be unique

in the sense of the probability law; see, for instance, Liptser and Shiriyayev [1977].

Consider now a nonnegative function ϕ on the reals which is even, $C^2(\mathbb{R})$, uniformly convex in the sense that

$$(2.3) \quad 0 < k \leq \ddot{\phi}(x) \leq K, \quad \text{all } x \in \mathbb{R}$$

for some positive constants k, K , with $\ddot{\phi}(x)$ decreasing on $x > 0$.

The control problem consists in finding a law $u^* \in \mathcal{U}$ that minimizes the "discounted" expected total cost

$$(2.4) \quad J(x; u) = E \int_0^\infty e^{-\alpha t} (|u_t(x)| + \phi(x_t)) dt$$

of starting at place x and using control u , over all $u \in \mathcal{U}$, $x \in \mathbb{R}$. Here E denotes expectation with respect to the probability measure P , $\alpha > 0$ is the "discount factor", $\phi(\cdot)$ is the running cost on the state and $|\cdot|$ is the cost of control.

3. THE EQUATION OF DYNAMIC PROGRAMMING

The method proceeds by constructing a solution to the Bellman equation of dynamic programming which satisfies certain growth and symmetry conditions. Introduce the function

$$(3.1) \quad a(p) = \min_{|u| \leq 1} (|u| + pu) = \begin{cases} 0 & , \quad |p| \leq 1 \\ 1 - |p| & , \quad |p| \geq 1. \end{cases}$$

The formal Bellman equation for this problem is

$$(3.2) \quad \alpha v = \frac{1}{2} v_{xx} + a(v_x) + \phi(x), \quad x \in \mathbb{R}.$$

We are looking for a positive number b and an even solution of

$$(3.2) \quad v(x) = O(x^2) \quad \text{as } |x| \rightarrow \infty, \text{ such that } v_x(b) = 1 \text{ and}$$

$$(3.3) \quad \alpha v = \frac{1}{2} v_{xx} + \phi(x), \quad 0 < v(x) < 1 \quad \text{on } 0 < x < b$$

$$(3.4) \quad \alpha v = \frac{1}{2} v_{xx} + 1 - v_x + \phi(x), \quad v_x(x) > 1 \quad \text{on } x > b.$$

A particular solution to the equation in (3.3) is given by the cost of "doing nothing" all the time. Indeed, consider the "naive" control law $u_t(x) \equiv 0$. The corresponding cost is

$$p(x) = E \int_0^\infty e^{-\alpha t} \phi(x + w_t) dt$$

and it becomes an easy exercise in Laplace transforms to verify that

$$(3.5) \quad p(x) = \frac{1}{\sqrt{2\alpha}} \left[e^{-x\sqrt{2\alpha}} \int_{-\infty}^x \phi(z) e^{z\sqrt{2\alpha}} dz + \int_x^{\infty} \phi(z) e^{-z\sqrt{2\alpha}} dz \right].$$

This function is even, has the growth of $\phi(\cdot)$ as $|x| \rightarrow \infty$ and satisfies the equation in (3.3) as is easily verified. To get the general solution of the latter, we add to $p(x)$ a solution

$$A_1 e^{x\sqrt{2\alpha}} + A_2 e^{-x\sqrt{2\alpha}}$$

of the homogeneous $\alpha v = \frac{1}{2} v_{xx}$. Since $v(\cdot)$ has to be even, and consequently $v_x(0) = 0$, $A_1 = A_2 = \frac{A}{2}$. So

$$(3.6) \quad v(x) = A \cosh(x\sqrt{2\alpha}) + p(x), \quad \text{on } 0 < x < b.$$

Condition $v_x(b) = 1$ then implies

$$(3.7) \quad A = \frac{p'(b) - 1}{\sqrt{2\alpha} \cdot \sinh(b\sqrt{2\alpha})}.$$

Similarly, a particular solution to (3.4) is obtained by considering the cost corresponding to the naive law $u_t(x) \equiv -1$ of pushing with full force to the left all the time:

$$q(x) \triangleq E \int_0^{\infty} e^{-\alpha t} (1 + \phi(x-t+w_t)) dt$$

and is easily verified that, if $\beta \triangleq \sqrt{1+2\alpha} - 1$,

$$(3.8) \quad q(x) = \frac{1}{\alpha} + \frac{1}{1+\beta} \left[e^{-\beta x} \int_{-\infty}^x \phi(z) e^{\beta z} dz + e^{(2+\beta)x} \int_x^{\infty} \phi(z) e^{-(2+\beta)z} dz \right],$$

solves the equation in (3.4) and has the growth of $\phi(\cdot)$ as $|x| \rightarrow \infty$. To get the general solution of (3.4) one has to add to $q(x)$ the general solution

$$B e^{-\beta x} + B_1 e^{(2+\beta)x}$$

of the corresponding homogeneous equation $\alpha v = \frac{1}{2} v_{xx} - v_x$ (note that $2 + \beta$ and $-\beta$ are the roots of the characteristic polynomial $s^2 - 2s - 2\alpha$). The growth condition implies $B_1 = 0$, so

$$(3.9) \quad v(x) = B e^{-\beta x} + q(x), \quad \text{on } x > b,$$

where

$$(3.10) \quad B = \frac{q'(b) - 1}{\beta} e^{\beta b}$$

because of $v_x(b) = 1$. Matching the values of $v(\cdot)$ from the two sides at $x = b$ gives the equation for the switching point b :

$$(3.11) \quad \tanh(b\sqrt{2\alpha}) = - \frac{1}{\sqrt{2\alpha}} \frac{p'(b) - 1}{\frac{q'(b) - 1}{\beta} + q(b) - p(b)},$$

$p(\cdot)$ and $q(\cdot)$ being the functions in (3.5), (3.8).

Proposition 3.1. There is exactly one positive solution b to equation (3.11).

Proof. A lot of simple calculus shows that $p'(x) - 1 = \frac{1}{\alpha} m_1(x)$

and $\frac{q'(x) - 1}{\beta} + q(x) - p(x) = \frac{\beta}{2\alpha} m_2(x)$, where

$$m_1(x) \triangleq \phi'(x) - \alpha + \frac{1}{2} \left\{ e^{x\sqrt{2\alpha}} \int_x^\infty \phi''(z) e^{-z\sqrt{2\alpha}} dz - e^{-x\sqrt{2\alpha}} \int_{-\infty}^x \phi''(z) e^{z\sqrt{2\alpha}} dz \right\}$$

$$m_2(x) \triangleq \phi'(x) - \alpha + e^{(2+\beta)x} \int_x^\infty \phi''(z) e^{-(2+\beta)z} dz - \frac{\sqrt{2\alpha}}{\beta} \left\{ e^{x\sqrt{2\alpha}} \int_x^\infty \phi''(z) e^{-z\sqrt{2\alpha}} dz + e^{-x\sqrt{2\alpha}} \int_{-\infty}^x \phi''(z) e^{z\sqrt{2\alpha}} dz \right\}$$

so that equation (3.11) becomes: $\tanh(x\sqrt{2\alpha}) = \frac{\sqrt{2\alpha}}{\beta} m(x)$,

$$m(x) \triangleq -\frac{m_1(x)}{m_2(x)}.$$

Note that $m_1(0) = -\alpha < 0$ and that $m_1(x)$ is strictly increasing to infinity as $x \rightarrow \infty$, since

$$m_1'(x) = \sqrt{\frac{\alpha}{2}} \left(e^{x\sqrt{2\alpha}} \int_x^\infty \phi''(z) e^{-z\sqrt{2\alpha}} dz + e^{-x\sqrt{2\alpha}} \int_{-\infty}^x \phi''(z) e^{z\sqrt{2\alpha}} dz \right) \geq k > 0.$$

Thus, there exists a unique number $b_1 > 0$, such that $m_1(b_1) = 0$.

On the other hand, since $\beta < \sqrt{2\alpha} < 2 + \beta$, $m_1(x) > m_2(x)$ and

$$m_2'(x) = \frac{\alpha}{\beta} \left[2 \int_x^\infty \phi''(z) e^{-(2+\beta)(z-x)} dz + \int_0^\infty (\phi''(z-x) - \phi''(z+x)) e^{-z\sqrt{2\alpha}} dz \right] \geq k > 0$$

on $x > 0$, by the assumption of decreasing curvature. So there exists a unique number $b_2 > b_1$, such that $m_2(b_2) = 0$. Now the function $m(x)$ is negative on $(0, b_1)$ and (b_2, ∞) , is equal to zero at b_1 , and increases monotonically to infinity on (b_1, b_2) as $x \uparrow b_2$.

Consequently, there exists a unique $b \in (b_1, b_2)$ such that:
 $\tanh(b\sqrt{2\alpha}) = \frac{\sqrt{2\alpha}}{\beta} m(b)$, q.e.d.

Once b has been thus determined, one constructs the function

$$\begin{aligned} (3.12) \quad v(x) &= \frac{1 - p'(b)}{\sqrt{2\alpha} \cdot \sinh(b\sqrt{2\alpha})} \cosh(x\sqrt{2\alpha}) + p(x) ; \quad 0 \leq x \leq b \\ &= \frac{q'(b) - 1}{\beta} e^{-\beta(x-b)} + q(x) ; \quad x > b \\ &= v(-x) ; \quad x < 0 \end{aligned}$$

in accordance with (3.6), (3.7), (3.9), (3.10), where $p(x)$ and $q(x)$ are again the functions in (3.5) and (3.8). The function $v(\cdot)$ in (3.12) satisfies equations (3.3) and (3.4) on (a, b) and (b, ∞) respectively, as well as $v_x(b) = 1$, by construction, and $v(b_+) = v(b_-)$, by (3.11).

It remains to prove that $v(\cdot)$ solves the Bellman equation (3.2). Suffices to prove: $0 \leq v_x(x) \leq 1$, on $[0, b]$ and

$v_x(x) \geq 1$, on $[b, \infty)$, while in turn this is an easy corollary of convexity:

Proposition 3.2. The function in (3.12) is convex: $v_{xx}(x) \geq 0$, $x \in \mathbb{R}$.

Proof. A bit of algebra shows that, on $0 \leq x \leq b$,

$$\begin{aligned} \frac{1}{2} v_{xx}(x) = \alpha v(x) - \phi(x) &\geq \frac{\beta}{2\alpha} \left[\phi'(b) + e^{(2+\beta)b} \int_b^\infty \phi''(z) e^{-(2+\beta)z} dz \right] \\ &+ (\alpha p(x) - \phi(x)) - (\alpha p(b) - \phi(b)). \end{aligned}$$

Note that: $\alpha p'(x) - \phi'(x) = \frac{1}{2} \int_0^\infty (\phi''(z+x) - \phi''(z-x)) e^{-z\sqrt{2\alpha}} dz \leq 0$, by the decreasing second derivative assumption. Therefore, $\alpha p(x) - \phi(x) \geq \alpha p(b) - \phi(b)$ and

$$(3.13) \quad v_{xx}(x) \geq \frac{\beta k(b+1)}{\alpha}, \quad 0 \leq x \leq b.$$

By continuity of $v_{xx}(\cdot)$; $v_{xx}(x) > 0$ on $(b, b+\epsilon)$, $\epsilon > 0$ sufficiently small. On the other hand, if $w = v_{xx}$:

$$w_{xx} - 2w_x - 2\alpha w = -2\phi''(x) \leq 0, \quad \text{on } (b, \infty).$$

By the maximum principle (Friedman [1964], p. 53, Theorem 18), $v_{xx}(\cdot)$ cannot have a negative minimum on (b, ∞) . However, on this interval,

$$v_{xx}(x) = \beta(q'(b)-1)e^{-\beta(x-b)} + q''(x) \geq \beta(q'(b)-1)e^{-\beta(x-b)} + \frac{k}{\alpha},$$

since

$$q''(x) = \frac{1}{1+\beta} \left[e^{(2+\beta)x} \int_x^{\alpha} \phi''(z) e^{-(2+\beta)z} dz + e^{-\beta x} \int_{-\infty}^x \phi''(z) e^{\beta z} dz \right].$$

Therefore, $v_{xx}(x) > 0$ for x sufficiently large, so if $v_{xx}(\bar{x}) < 0$, some $\bar{x} \in (b, \infty)$, $v_{xx}(\cdot)$ would have a negative minimum there, contradicting the maximum principle. Therefore $v_{xx}(x) \geq 0$, also on $[b+\epsilon, \infty)$, q.e.d.

Special Case: In the special case $\phi(x) = x^2$, we have

$$p(x) = \frac{x^2}{\alpha} + \frac{1}{\alpha^2}, \quad q(x) = \frac{x^2}{\alpha} - \frac{2}{\alpha}x + \frac{\alpha^2 + \alpha + 2}{\alpha^2},$$

$m_1(x) = x - \frac{\alpha}{2}$, $m_2(x) = x - (\frac{\alpha}{2} + \frac{1}{\alpha})$, so $\frac{\alpha}{2} < b < \frac{\alpha}{2} + \frac{1}{\alpha}$. It can be shown that $v_{xx}(x) \geq \frac{2(\alpha-\beta)}{\alpha^2} > 0$, any $x \in \mathbb{R}$, in this case.

4. THE OPTIMAL LAW

Let us prove that $v(\cdot)$ is the value function of the control problem, i.e. an (attainable) lower bound on the expected discounted total cost, and try to discern the law that achieves this infimum. Consider any admissible law $u \in \mathcal{U}$ along with the corresponding state process (x_t^u) , solving equation (2.1)-(2.2) in the weak sense. We introduce the process

$$(4.1) \quad V_t^u \triangleq v(x_t^u)e^{-\alpha t}$$

and note that

$$Ev(x_t^u) \leq Ev(|x| + t + |w_t|) = O(t^2) \quad \text{as } t \rightarrow \infty, \text{ so } \lim_{t \rightarrow \infty} EV_t^u = 0,$$

any $\alpha > 0$, $x \in \mathbb{R}$, $u \in \mathcal{U}$.

Applying Itô's rule to (4.1) we get

$$\begin{aligned} V_T^u = v(x) &+ \int_0^T e^{-\alpha t} (-\alpha v(x_t^u) + \frac{1}{2} v_{xx}(x_t^u) + u_t v_x(x_t^u) + |u_t| + \\ &+ \phi(x_t^u)) dt - \int_0^T e^{-\alpha t} (|u_t(x^u)| + \phi(x_t^u)) dt + \int_0^T e^{-\alpha t} v_x(x_t^u) dw_t. \end{aligned}$$

Since $v(\cdot)$ satisfies equation (3.2), the first integrand is nonnegative. Taking expectations and then passing to the limit as $T \rightarrow \infty$ we get

$$(4.2) \quad J(x; u) = E \int_0^\infty e^{-\alpha t} (|u_t(x^u)| + \phi(x_t^u)) dt \geq v(x); \quad u \in \mathcal{U}, x \in \mathbb{R}.$$

Consider now the law $u^* \in \mathcal{U}$

$$(1.1) \quad u_t^*(x) = \begin{cases} -1, & x_t > b \\ 0, & |x_t| \leq b \\ 1, & x_t < -b \end{cases}$$

obtained through the minimization: $|u_t^*(x)| + v_x(x_t) \cdot u_t^*(x) = a(v_x(x_t))$.
The corresponding state process (x_t^*) satisfies

$$(4.4) \quad \begin{aligned} dx_t^* &= u_t^*(x^*)dt + dw_t; \quad t \geq 0 \\ x_0^* &= x \end{aligned}$$

on an appropriate probability space. Although no explicit use is made of this fact in the present context, we mention that (4.4) is strongly solvable for x^* as a causal functional of w , because $u_t^*(x)$ in (4.3) is instantaneous, bounded and measurable; see Zvonkin [1974]. Then the inequalities above hold as equalities, and

$$(4.5) \quad J(x; u^*) = v(x); \quad x \in \mathbb{R}.$$

From (4.2), (4.5), $v(x)$ is a lower bound on the performance (2.4), and is achieved by the process (x_t^*) . In other words, $u_t^*(x)$ is optimal.

REFERENCES

- V.E. BENES[✓], L.A. SHEPP, H.S. WITSENHAUSEN [1979]: Some solvable stochastic control problems, preprint.
- H.J. KUSHNER [1967]: Optimal discounted stochastic control for diffusion processes. SIAM J. Control 5(4), pp. 520-531.
- R.S. LIPTSER, A.N. SHIRYAYEV [1977]: Statistics of Random Processes, Vol. I, (Engl. Transl.), Springer-Verlag, Berlin.
- A.K. ZVONKIN [1974]: A transformation of the phase space of a diffusion process that removes the drift, Math. USSR(Sbornik), (Engl. Transl.), 22(1), pp. 129-149.
- A. FRIEDMAN [1964]: Partial Differential Equations of Parabolic Type, Prentice-Hall, Englewood Cliffs, N.J.